



6016VVR

18% Show Goat Pellet

Medicated

A goat started feed formulated for starting and growing goats; medicated for the prevention of coccidiosis cause by Eimeria crandallis, Eimeria christenseni and Eimeria ninakohlyakimovae.

Active Drug Ingredient

Monensin.....20g/ton

Guaranteed Analysis

Crude Protein (Min).....18%
(includes not more than 0.9% equivalent crude protein from non-protein nitrogen)
Crude Fat (Min.).....5.0%
Crude Fiber (Max.).....12.8%
Calcium (Ca) (Min.).....0.60%
Calcium (Ca) (Max.).....1.00%
Phosphorus (P) (Min.).....0.45%
Salt (NaCl) (Min.).....0.25%
Salt (NaCl) (Max.).....0.75%
Copper (Cu) (Min.).....30ppm
Selenium (Se) (Min.).....0.50 ppm
Vitamin A (Min.).....15,000 IU/lb

Ingredients

Wheat Middlings, Corn Distillers Grains Dried, Dehydrated Alfalfa Meal, Corn, Flaxseed Meal, Corn Starch, Dehulled Soybean Meal, Soybean Hulls, Wheat Flour, Corn Gluten Meal, Miconcapsulated Dry Active Saccharomyces Cerevisiae, Molasses, Salt, Calcium Carbonate, Ammonium Chloride, Potassium Sulfate, Magnesium Sulfate, Animal Fat, Dried Extracted Aspergillus Fermentation Solubles, Saccharomyces Cerevisiae, Aspergillus Oryzae, Yucca Schidigera, Aspergillus Niger, Bacillus Subtilis, Lactobacillus, Acidophilus, Enterococcus Faecium, Bifobacterium Longum, Bifobacterium Thermophilum, Dried Aspergillus Oryzae Fermentation Products, Vitamin A Supplement, Vitamin E Supplement, Vitamin D-3 Supplement, Niacin, Vitamin B-12 Supplement, Riboflavin, d-Calcium Pantothenate, Folic Acid, Pyroxide Hydrochloride, Mena-dione Dimethylpyrimidinol Bisulfate, Ferrous Sulfate, Manganese Sulfate, Copper Sulfate, Ethylene Diamine Dihydriodide, Colbalt Sulfate, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper Amino Acid Complex, Cobalt Glucoheptonate, Selenium Yeast, and Sodium Selenite.

Caution: Do not allow horses or other equines access to feeds containing monensin. Ingestion of monensin by equines has been fatal. Feeding undiluted or mixing errors resulting in high concentrations of monensin has been fatal in cattle and could be fatal in goats. Monensin- medicated feed is for use in cattle or goats. Consumption of monensin in unapproved species may result in toxic reaction. Do not feed to lactating goats.

CONTAINS COPPER - DO NOT FEED TO SHEEP!

Feeding Directions

Depending on animal weight, body condition, and forage quality, feed 0.25 to 3lbs of 6618 VV along with good quality grass/legume hay. Feed as a creep feed from birth to 30 lbs or from birth to show. Provide clean, fresh water at all times.

Manufactured for

Brubaker Grain

307 W Center St, Farmersville, OH 45325

Net Weight - 50 LBS. (22.7Kg.) - BULK - Shown on Invoice